

The Topological Structure of Nieh-Yan Form and Chiral Anomaly in Spaces with Torsion

Sheng Li*

CCAST World Lab, Academia Sinica, Beijing 100080, P. R. China

Institute of Theoretical Physics, Academia Sinica, P. O. Box 2735, Beijing 100080, P. R. China

Abstract

The topological structure of the Nieh-Yan form in 4-dimensional manifold is given by making use of the decomposition of spin connection. The case of the generalized Nieh-Yan form on 2^d -dimensional manifold is discussed with an example of 8-dimensional case studied in detail. The chiral anomaly with nonvanishing torsion is studied also. The further contributions from torsional part to chiral anomaly are found coming from the zeroes of some fields under pure gauge condition.

PACS numbers: 02.40.-k, 11.15.-q, 11.30.Rd, 04.62.+v

Typeset using REVTeX

*lisheng@itp.ac.cn

I. INTRODUCTION

Torsion might be the most unusual field in physics. Though it has been under investigation for more than two thirds of a century, there is no general agreement on its mathematical formulation nor on its physical significance. Vast funds of work on torsion had been done by many physicists (see [1–11] for examples) since it was proposed by E. Cartan [12] in 1920s.

Consider a compact manifold M with metric $g_{\mu\nu}$. There are two dynamically independent one-forms: the connection $\omega^{ab} = \omega_{\mu}^{ab} dx^{\mu}$ and vielbein $e^a = e_{\mu}^a dx^{\mu}$. The curvature and torsion 2-forms are defined by

$$R^{ab} = d\omega^{ab} - \omega^{ac} \wedge \omega^{cb} \quad (1)$$

$$T^a = de^a - \omega^{ab} \wedge e^b. \quad (2)$$

In geometry, ω^{ab} and e^a reflect the affine and metric properties of M . While in physics, curvature and torsion may be related to energy momentum tensor and spin current respectively.

Nieh and Yan [13] firstly gave the 4-dimensional torsional invariant 4-form as

$$N = T^a \wedge T^a - R^{ab} \wedge e^a \wedge e^b. \quad (3)$$

This is the only nontrivial locally exact 4-form which vanishes in the absence of torsion and is clearly independent of the Pontryagin and Euler densities. In any local patch where the vielbein is well defined, N can be written as

$$N = d(e^a \wedge T^a) \quad (4)$$

and therefore is locally exact. The 3-form $e^a \wedge T^a$ is a Chern-Simons-like form that can be used as a Lagrangian for the dreibein in three dimensions. The dual of this 3-form in four dimensions is also known as the totally antisymmetric part of the torsion and is sometimes also referred to as H-torsion,

$$e^a \wedge T^a \wedge dx^{\rho} = \varepsilon^{\mu\nu\lambda\rho} T_{\mu\nu\lambda} d^4x. \quad (5)$$

This component of the torsion tensor is the one that couples to the spin 1/2 fields [7].

Recently, there have been some discussions on the question of further contributions to the Chiral anomaly in the presence of space-time with torsion [5–11]. It is shown the Nieh-Yan form does contribute to the chiral anomaly for massive field. This further anomaly term is associated with vacuum polarization diagrams with two external axial torsion vertices, rather than with the usual triangle diagrams [11].

In this paper, by making use of the decomposition theory of $SO(N)$ spin connection (see [16] for example) reviewed in section 2, we give the topological structure of Nieh-Yan form in 4-dimensional manifold in section 3. The Nieh-Yan number is found to be the sum of indices of some field ϕ at its zeroes. The Hopf indices and Brouwer degrees of ϕ label the local properties of the Nieh-Yan form. We also present the relationship between Nieh-Yan number and winding number. In section 4, a general discussion on the cases of generalized Nieh-Yan form on 2^d -dimensional manifold is given with a elaborate study on

8-dimensional case as an example. By applying the results obtained in this paper and the results obtained by Duan and Fu in ref. [21] to chiral anomaly, it is found under pure gauge condition, contributions by chiral anomaly come only from the zeroes of some field $\tilde{\phi}_{L(R)}$ and ϕ in section 5. At last, we give a short conclusion in section 6. One should notice that the main result of this paper is based on reduction to contribution by singular points, i.e. those where ordinary formulas do not hold.

II. THE DECOMPOSITION THEORY OF SPIN CONNECTION

In this section, we give a short review of the decomposition theory of $SO(N)$ which is a useful tool in the discussion of the topological structure of Nieh-Yan form in the later sections.

A smooth vector field φ^a ($a = 1, 2, \dots, N$) can be found on the base manifold $\mathbf{M}$ (a section of a vector bundle over $\mathbf{M}$). We define a unit vector n on $\mathbf{M}$ as

$$n^a = \varphi^a / \|\varphi\| \quad a = 1, 2, \dots, N \quad (6)$$

$$\|\varphi\| = \sqrt{\varphi^a \varphi_a},$$

in which the superscript “ a ” is the local orthonormal frame index. In fact n is identified as a section of the sphere bundle over $\mathbf{M}$ (or a partial section of the vector bundle over $\mathbf{M}$). We see that the zeroes of ϕ are just the singular points of n .

Let the N -dimensional Dirac matrix γ_a ($a = 1, 2, \dots, N$) be the basis of the Clifford algebra which satisfies

$$\gamma_a \gamma_b + \gamma_b \gamma_a = 2\delta_{ab}. \quad (7)$$

A unit vector field n on $\mathbf{M}$ can be expressed as a vector of Clifford Algebra

$$n = n^a \gamma_a, \quad (8)$$

The spin connection 1-form and curvature 2-form are represented as Clifford-algebra-valued differential forms respectively

$$\omega = \frac{1}{2} \omega^{ab} I_{ab} \quad F = \frac{1}{2} F^{ab} I_{ab}, \quad (9)$$

in which I_{ab} is the generator of the spin representation of the group $SO(N)$

$$I_{ab} = \frac{1}{4} [\gamma_a, \gamma_b] = \frac{1}{4} (\gamma_a \gamma_b - \gamma_b \gamma_a). \quad (10)$$

The covariant derivative 1-form of n^a can be represented in terms of n and ω as

$$Dn = dn - [\omega, n], \quad (11)$$

and curvature 2-form is

$$F = d\omega - \omega \wedge \omega. \quad (12)$$

Arbitrary $U \in Spin(N)$, which satisfies

$$UU^\dagger = U^\dagger U = I, \quad (13)$$

is an even-versor [14]. The induced ‘spinorial’ transformation by U to the basis γ_i of the Clifford algebra give N orthonormal vectors u_i [15] via

$$u_i := U\gamma_i U^\dagger = u_i^a \gamma_a, \quad (14)$$

where u_i^a is the coefficient of u_i in the representation of Clifford algebra. From the relationship between U and u_i^a , we see that u_i has the same singular points with respect to different “ i ”. By (14), it is easy to verify that u_i satisfy

$$u_i u_j + u_j u_i = 2\delta_{ij}, \quad i, j = 1, 2, \dots, N. \quad (15)$$

From (11) we know that the covariant derivative 1-form of u_i is

$$Du_i = du_i - [\omega, u_i]. \quad (16)$$

There exists the following formula for a Clifford Algebra r -vector A [14]

$$u_i A u_i = (-1)^r (N - 2r) A. \quad (17)$$

For ω is a Clifford Algebra 2-vector and using (17), the spin connection ω can be decomposed by N orthonormal vectors u_i as

$$\omega = \frac{1}{4} (du_i u_i - Du_i u_i) \quad (18)$$

or

$$\omega^{ab} = du_i^a u_i^b - Du_i^a u_i^b. \quad (19)$$

It can be proved that the general decomposition formula (18) has global property and is independent of the choice of the local coordinates [16].

By choosing the gauge condition

$$Du_i = 0, \quad (20)$$

we can define a generalized pseudo-flat spin connection as

$$\omega_0 = \frac{1}{4} du_i u_i. \quad (21)$$

Suppose there exist l singular points z_i ($i = 1, 2, \dots, l$) in the orthonormal vectors u_j . One can easily prove that at the normal points of u_j

$$F(\omega_0) = 0 \quad \text{when } x \neq z_i. \quad (22)$$

For the derivative of u_i at the singular points z_i is undefined, formula (22) is invalid at z_i . Hence, the curvature under the gauge condition (20) is a generalized function

$$F \begin{cases} = 0 \\ \neq 0 \end{cases} \quad \text{when } \begin{cases} x \neq z_i, \\ x = z_i. \end{cases} \quad (23)$$

This is why we call ω_0 the pseudo-flat spin connection. In fact, the gauge condition (20) is the pure gauge condition supported by the fact that one can always find a frame which is locally flat.

III. TOPOLOGICAL STRUCTURE OF NIEH-YAN FORM IN 4-DIMENSIONAL MANIFOLD

One of the properties of topological invariant is that it is independent of connection. Therefore, we choose the pseudo-flat connection

$$\omega^{ab} = du_i^a u_i^b \quad (24)$$

to make the calculation easier. Under this gauge condition, there must exist singularity points on the manifold if the topology of this manifold is nontrivial. In fact, what we have done is to choose a frame which is locally flat. The choice of frame does not change the topology of the manifold. For the singular property of the pseudo-flat connection, the contribution by the zeros of some field at its singular points will give the topological characteristic. It is a useful method to choose a locally flat frame when dealing with the topological properties of manifolds, see for example [22].

Using the pseudo-flat spin connection, we can rewrite torsion as

$$T^a = De^a = D(e_i u_i^a) = de_i u_i^a \quad (25)$$

and Nieh-Yan form

$$N = d(e^a \wedge T^a) = de_i \wedge de_i, \quad (26)$$

where e_i are the projection of vielbein onto the basis

$$e_i = e^a u_i^a. \quad (27)$$

In quaternionic representation, a four vector $\vec{\phi} = (\phi_1, \phi_2, \phi_3, \phi_4)$ is written as

$$\phi = \phi_1 i + \phi_2 j + \phi_3 k + \phi_4 \quad \phi^* = -\phi_1 i - \phi_2 j - \phi_3 k + \phi_4 \quad (28)$$

and the vielbein projection

$$e = e_1 i + e_2 j + e_3 k + e_4, \quad (29)$$

where $(1, i, j, k)$ is the basis of quaternion satisfying $i^2 = -1$, $ij = k$, etc. To give a frame that is flat locally, we can find some ϕ , which make the vielbein expressed in the form

$$e = \frac{l}{||\phi||^2} \phi d\phi^*, \quad (30)$$

where the constant l has dimension of length. The choice of local flat metric is coincide with the pseudo-flat connection chosen, which makes singular points exist for the non-trivial topological property of the manifold..

It's well known that the quaternionic representation can be expressed in terms of the Clifford algebra as

$$\phi = \phi_i s^i \quad i = 1, 2, 3, 4, \quad (31)$$

where

$$s = (i\vec{\sigma}, I), \quad s^\dagger = (-i\vec{\sigma}, I) \quad (32)$$

and

$$e = e_i s^i. \quad (33)$$

The vielbein projection can be rewritten as

$$e = \frac{l}{||\phi||^2} \phi d\phi^\dagger. \quad (34)$$

Then we get the Nieh-Yan form as

$$N = de_i \wedge de_i = \frac{1}{2} Tr(de \wedge de). \quad (35)$$

By making use of the relationship

$$\varepsilon^{ijkl} = \frac{1}{2} Tr(s^i s^j s^k s^l), \quad (36)$$

the Nieh-Yan form can be expressed in terms of a unit vector $n_i = \frac{\phi_i}{||\phi||}$ as

$$\begin{aligned} N &= \frac{l^2}{2} Tr(dn \wedge dn \wedge dn \wedge dn) \\ &= l^2 \varepsilon^{ijkl} dn_i \wedge dn_j \wedge dn_k \wedge dn_l. \end{aligned} \quad (37)$$

The derivative of n_a can be deduced as

$$dn_i = \frac{d\phi_i}{||\phi||} - \phi_i d\left(\frac{1}{||\phi||}\right). \quad (38)$$

Substituting it into (37), we have the expression of N on $S(\mathbf{M})$

$$N = l^2 \varepsilon^{ijkl} d\left(\frac{\phi_i}{||\phi||^4}\right) d\phi_j \wedge d\phi_k \wedge d\phi_l. \quad (39)$$

Using

$$\frac{\phi_i}{||\phi||^4} = -\frac{1}{2} \frac{\partial}{\partial \phi_i} \left(\frac{1}{||\phi||^2} \right), \quad (40)$$

equation (39) is

$$\begin{aligned} N &= -\frac{l^2}{2} \varepsilon^{ijkl} \frac{\partial}{\partial \phi_m} \frac{\partial}{\partial \phi_i} \left(\frac{1}{||\phi||^2} \right) \\ &\quad \times \frac{\partial \phi_m}{\partial x^\mu} \frac{\partial \phi_j}{\partial x^\nu} \frac{\partial \phi_k}{\partial x^\lambda} \frac{\partial \phi_l}{\partial x^\rho} \frac{\epsilon^{\mu\nu\lambda\rho}}{\sqrt{g}} \sqrt{g} d^4x, \end{aligned} \quad (41)$$

where $g = \det(g_{\mu\nu})$, $g_{\mu\nu}$ is the metric tensor of $\mathbf{M}$. Define the Jacobian $D(\phi/x)$ as

$$\varepsilon^{ijkl} D(\phi/x) = \epsilon^{\mu\nu\lambda\rho} \frac{\partial\phi_i}{\partial x^\mu} \frac{\partial\phi_j}{\partial x^\nu} \frac{\partial\phi_k}{\partial x^\lambda} \frac{\partial\phi_l}{\partial x^\rho}. \quad (42)$$

Noticing

$$\varepsilon^{ijkl} \varepsilon^{mjkl} = 6\delta^{im}, \quad (43)$$

we get

$$N = -3l^2 \frac{\partial^2}{\partial\phi_i \partial\phi_i} \left(\frac{1}{\|\phi\|^2} \right) D\left(\frac{\phi}{x}\right) d^4x. \quad (44)$$

The general Green's-function formula [17] in ϕ space is

$$\frac{\partial^2}{\partial\phi_i \partial\phi_i} \left(\frac{1}{\|\phi\|^2} \right) = -4\pi^2 \delta(\phi), \quad (45)$$

We obtain the new formulation of Nieh-Yan form in terms of δ function $\delta(\phi)$

$$N = 12\pi^2 l^2 \delta(\phi) D(\phi/x) d^4x. \quad (46)$$

Suppose $\phi(x)$ has l isolated zeroes on $\mathbf{M}$ and let the i th zero be z_i , it is well known from the ordinary theory of the δ -function [18] that

$$\delta(\phi) = \sum_{i=1}^l \frac{\beta_i \delta(x - z_i)}{D(\phi/x)|_{x=z_i}}. \quad (47)$$

Then one obtains

$$\delta(\phi) D\left(\frac{\phi}{x}\right) = \sum_{i=1}^l \beta_i \eta_i \delta(x - z_i), \quad (48)$$

where β_i is the positive integer (the Hopf index of the i th zero) and η_i the Brouwer degree [19,20]

$$\eta_i = \text{sgn} D(\phi/x)|_{x=z_i} = \pm 1. \quad (49)$$

From the above deduction the following topological structure is obtained:

$$n_{N-Y} = 12\pi^2 l^2 \delta(\phi) D\left(\frac{\phi}{x}\right) d^4x = 12\pi^2 l^2 \sum_{i=1}^l \beta_i \eta_i \delta(x - z_i) d^4x, \quad (50)$$

which means that the local structure of N is labeled by the Brouwer degrees and Hopf indices, which are topological invariants. Therefore the Nieh-Yan number n_{N-Y} can be represented as

$$n_{N-Y} = \int_{\mathbf{M}} N = 12\pi^2 l^2 \sum_{i=1}^l \beta_i \eta_i. \quad (51)$$

On another hand, we can decompose $\mathbf{M}$ as

$$\mathbf{M} = \sum_i \mathbf{M}_i, \quad (52)$$

so that $\mathbf{M}_i$ includes only the i th singularity point z_i of $n(x)$. Then we get

$$\begin{aligned} n_{N-Y} &= \sum_i \int_{\mathbf{M}_i} l^2 \varepsilon^{ijkl} dn_i \wedge dn_j \wedge dn_k \wedge dn_l \\ &= \sum_i \oint_{\partial \mathbf{M}_i} l^2 \varepsilon^{ijkl} n_i \wedge dn_j \wedge dn_k \wedge dn_l, \end{aligned} \quad (53)$$

where $\partial \mathbf{M}_i$ is the boundary of $\mathbf{M}_i$. Equation (53) is another definition of the winding number $W(\phi, z_i)$ of the surface $\partial \mathbf{M}_i$ and the mapping $\phi(x)$ [23]

$$W(\phi, z_i) = \frac{1}{12\pi^2} \oint_{\partial \mathbf{M}_i} \varepsilon^{ijkl} n_i \wedge dn_j \wedge dn_k \wedge dn_l = \beta_i \eta_i. \quad (54)$$

Then the Nieh-Yan number n_{N-Y} can further be expressed in terms of winding numbers

$$n_{N-Y} = 12\pi^2 l^2 \sum_{i=1}^l W(\phi, z_i). \quad (55)$$

The sum of the winding numbers can be interpreted or, indeed, defined as the degree of the mapping $\phi(x)$ onto $\mathbf{M}$. By (46) and (53), we have

$$\begin{aligned} \sum_{i=1}^l W(\phi, z_i) &= \int_{\mathbf{M}} \delta(\phi) D\left(\frac{\phi}{x}\right) d^4x \\ &= \deg \phi \int_{\phi(\mathbf{M})} \delta(\phi) d^4\phi \\ &= \deg \phi. \end{aligned} \quad (56)$$

Therefore, we get the Nieh-Yan number at last

$$n_{N-Y} = 12\pi^2 l^2 \sum_{i=1}^l W(\phi, z_i) = 12\pi^2 l^2 \deg \phi. \quad (57)$$

From (6), we know that the zeroes of ϕ are just the singularities of n . Here (51) says that the sum of the indices of the singular points of n , or of the zeroes of ϕ , is Nieh-Yan number. One should notice that the vector field ϕ is not a section of the tangent bundle of $\mathbf{M}$, i.e. ϕ is not a Riemannian vector field. Therefore formula (57) does not concern with the Poincaré-Hopf theorem which connects the Euler characteristics with the zeros of a Riemannian vector field.

IV. HIGHER DIMENSIONAL CASE

In 2^d -dimensional case, we can get similar results. Under the pure gauge condition, the nonvanishing term of the generalized Nieh-Yan form is

$$(de_i \wedge de_i)^{2^{d-2}}. \quad (58)$$

The computation above is also valid in this case. The topological structure of generalized Nieh-Yan form is constitute by delta function of some ϕ . The Hopf indices and Brouwer degree labeled the local structure of Nieh-Yan form. The Nieh-Yan number is some constant number times of the degree or winding numbers of ϕ .

$$N \sim \delta(\phi) D\left(\frac{\phi}{x}\right) d^{2^d} x = \sum_i \beta_i \eta_i \delta(x - z_i) d^{2^d} x. \quad (59)$$

In the following, we take only 8-dimensional case as an example.

In 8-dimensional compact manifold, under the gauge condition (20), the generalized Nieh-Yan form is expressed simply as

$$N = d(e_i \wedge de_i \wedge de_j \wedge de_j). \quad (60)$$

And the vielbein is expressed locally in terms of the octonions under Clifford algebra representation

$$e = e_i s^i = \frac{l}{\|\phi\|^2} \phi d\phi^\dagger, \quad (61)$$

where

$$\phi = \phi_i s^i \quad i = 1, 2, \dots, 8.$$

in which s^i is the basis of octonions. Substituting (61) into (60), we get the Nieh-Yan form of 8-d in terms of unit vector $n_i = \frac{\phi_i}{\|\phi\|}$ as

$$N = \frac{l^4}{5} \varepsilon^{i_1 i_2 \dots i_8} dn_{i_1} \wedge dn_{i_2} \wedge \dots \wedge dn_{i_8}. \quad (62)$$

Analogously to 4-dimensional case, using the general Green's-function formula [17] in 8-dimensional ϕ space, the generalized Nieh-Yan form is expressed as

$$\begin{aligned} N &= 1008 l^4 A(S^7) \delta(\phi) D\left(\frac{\phi}{x}\right) d^8 x \\ &= 336 \pi^4 l^4 \delta(\phi) D\left(\frac{\phi}{x}\right) d^8 x, \end{aligned} \quad (63)$$

which can be rewritten in terms of Hopf indices and Brouwer degree as

$$N = 336 \pi^4 l^4 \sum_i \beta_i \eta_i \delta(x - z_i) d^8 x. \quad (64)$$

The Nieh-Yan number in 8-dimensional is

$$n_{N-Y} = 336 \pi^4 l^4 \deg \phi = 336 \pi^4 l^4 \sum_i W(\phi, z_i). \quad (65)$$

Similar results can be obtained for higher 2^d -dimensional case.

V. CHIRAL ANOMALY ON SPACES WITH TORSION

Consider a massive Dirac spinor on a curved background with torsion. The action is

$$S = \int \frac{1}{2} (d^4x e \bar{\psi} \not{\nabla} \psi + h.c.) + m \bar{\psi} \psi, \quad (66)$$

where the Dirac operator is given by

$$\not{\nabla} = e^{\mu a} \gamma_a D_\mu. \quad (67)$$

This action is invariant under rigid chiral transformations

$$\psi' \rightarrow e^{i\varepsilon \gamma_5} \psi, \quad (68)$$

where ε is a real constant parameter. This symmetry leads to the classical conservation law

$$\partial_\mu J_5^\mu = 0, \quad (69)$$

in which

$$J_5^\mu = e e^{\mu a} \bar{\psi} \gamma_a \gamma_5 \psi. \quad (70)$$

The chiral anomaly when torsion is present is given by [5,11,6,8]

$$\partial_\mu < J_5^\mu > = A(x), \quad (71)$$

with

$$A(x) = \frac{1}{8\pi^2} * [R^{ab} \wedge R^{ab} + \frac{2}{l^2} (T^a \wedge T^a - R^{ab} \wedge e^a \wedge e^b)]. \quad (72)$$

The constant l is called the radius of the universe and is related to the cosmological constant ($|\Lambda| = l^{-2}$).

Using the results obtained in section 3, we can rewritten the chiral anomaly as

$$\partial_\mu < J_5^\mu > = \frac{1}{8\pi^2} * [R^{ab} \wedge R^{ab}] + 3\delta(\phi) D\left(\frac{\phi}{x}\right). \quad (73)$$

From the relationship

$$so(4) = su(2)_L \times su(2)_R, \quad (74)$$

the Pontryagin class of $SO(4)$ group can be expressed as the sum of the second Chern classes of the left and right $SU(2)_{L(R)}$ subgroup

$$Tr(R \wedge R) = Tr(R_{SU(2)_L} \wedge R_{SU(2)_L}) + Tr(R_{SU(2)_R} \wedge R_{SU(2)_R}). \quad (75)$$

By making use of the result of Duan and Fu [21]

$$\frac{1}{8\pi^2} Tr(R_{SU(2)} \wedge R_{SU(2)}) = \delta(\tilde{\phi}) D\left(\frac{\tilde{\phi}}{x}\right) d^4x, \quad (76)$$

where ϕ belongs to $Spin(3)$ corresponding to the group $SU(2)$. In Duan and Fu's paper, the pure gauge condition is used also to get the topological structure of the second Chern class for $SU(2)$ group. Now we get the chiral anomaly

$$\partial_\mu < J_5^\mu > = \delta(\tilde{\phi}_L) D\left(\frac{\tilde{\phi}_L}{x}\right) + \delta(\tilde{\phi}_R) D\left(\frac{\tilde{\phi}_R}{x}\right) + 3\delta(\phi) D\left(\frac{\phi}{x}\right), \quad (77)$$

in which $\tilde{\phi}_L$ and $\tilde{\phi}_R$ are some $Spin(3)_{L(R)}$ elements corresponding to the subgroups $SU(2)_{L(R)}$. Furthermore, by making use of the structure of delta function, the anomaly can be formulated more explicitly

$$\begin{aligned} \partial_\mu < J_5^\mu > &= \sum_i \delta(x - z_{Li}) \beta_{Li} \eta_{Li} + \sum_i \delta(x - z_{Ri}) \beta_{Ri} \eta_{Ri} + 3 \sum_i \delta(x - z_i) \beta_i \eta_i \\ &= \sum_i \delta(x - z_{Li}) W_{Li} + \sum_i \delta(x - z_{Ri}) W_{Ri} + 3 \sum_i \delta(x - z_i) W_i. \end{aligned} \quad (78)$$

where $\beta_{L(R)i}$ and $\eta_{L(R)i}$ are the Hopf indices and Brouwer degrees respectively corresponding to the zeroes of $\tilde{\phi}_{L(R)}$, and $W_{L(R)i}$ is the winding number of $\tilde{\phi}_{L(R)}$ at its i th zeroes. From (78), we see, under pure gauge condition, the chiral anomaly comes only from the zeroes of the fields $\tilde{\phi}_{LR}$ and ϕ , and their winding numbers account the quantity of chiral anomaly.

VI. CONCLUSION

In this paper, we have discussed the Nieh-Yan form by making use of the decomposition theory of spin connection. The Nieh-Yan form in 4-dimensional manifold is proved to take the delta function form under local flat gauge condition. The local topological structure of Nieh-Yan form is labeled by Hopf indices and Brouwer degrees of field ϕ , which is used to express the vielbein under local flat (or pure) gauge condition. The Nieh-Yan number is proved to be a constant number times the degree or winding number of ϕ .

A general discussion of the generalized Nieh-Yan form on 2^d -dimensional manifold is presented. From an example of 8-dimensional case, we find the topological structure of the generalized Nieh-Yan form is similar to the 4-dimensional case in terms of field ϕ under pure gauge condition. It is noticeable that the Clifford algebra can make the calculus more easier in these cases.

The topological anomaly with nonvanishing torsion is formulated explicitly under pure gauge condition. There are two kinds of contribution in the anomaly which comes from the topology of the manifold: the Pontryagin class of $SO(4)$ and the Nieh-Yan form. We have proved each of them to be a sum of delta function of some fields $\tilde{\phi}_{L(R)}$ and ϕ under pure gauge condition. It means the contribution comes only from the zeroes of $\tilde{\phi}_{L(R)}$ and ϕ . The degrees or winding numbers of $\tilde{\phi}_{L(R)}$ or ϕ give the quantity of contributions.

REFERENCES

- [1] T. Kibble *J. Math. Phys.* **2**, 212 (1961).
- [2] D. Sciama *Rev. Mod. Phys.* **36**, 463 (1964).
- [3] R. Hammond *Gen. Rel. Grav.* **22**, 451 (1990); **26**, 247 (1994).
- [4] Ramanand Jha *Gen. Rel. Grav.* **25**, 281 (1993); *Int. J. Mod. Phys.* **A9** 3595 (1994).
- [5] O. Chandia and J. Zanelli, *Phys. Rev.* **D55**, 7580 (1997), hep-th/9702025; *Phys. Rev.* **D58**, 045014 (1998) hep-th/9803034.
- [6] Y. N. Obukhov, E. W. Mielke, J. Budezies, and F. W. Hehl, *Found. Phys.* **27**, 1221 (1997) gr-qc/9702011.
- [7] Y. N. Obukhov, *Phys. Lett.* **B108**, 308 (1982); *Nucl. Phys.* **B212**, 237 (1983); *J. Phys.* **A16**, 3795 (1983).
- [8] E. W. Mielke, and D. Kreimer, *Chiral Anomaly in Contorted Spacetimes*, gr-qc/9704051.
- [9] D. Bellisai, *Nucl. Phys.* **B467**, 127 (1996).
- [10] H. Y. Guo, K. Wu and W. Zhang, to appear in *Comm. Theor. Phys.* hep-th/9805037;
- [11] C. Soo *Phys. Rev.* **D59**, 045006 (1999) hep-th/9805090.
- [12] È. Cartan *C. R. Acad. Sci.* **174**, 593 (1922).
- [13] H. T. Nieh and M. L. Yan, *J. math. Phys.* **23**, 373 (1982); *Ann. Phys.* **138**, 237 (1982).
- [14] D. Hestenes and G. Sobczyk, *Clifford Algebra to Geometric Calculus*. D. Reidel Publishing, (1984).
- [15] H. Boerner, *Representation of Groups*. North-Holland Publishing Company, (1963).
- [16] Y. S. Duan and S. Li, *Decomposition Theory of Spin Connection, Topological Structure of Gauss-Bonnet-Chern Topological Current and Morse Theory*, in *Jingshin Theoretical Physics Symposium in Honor of Professor Ta-You Wu*, World Scientific (1998).
- [17] I. M. Gelfand and G. E. Silov *Generalized Function*. Moscow, Nauka, Main Editorial for Mathematics and Physics (1959).
- [18] A. S. Schwarz, *Topology for Physicists*. Springer Verlag Press (1994).
- [19] B. A. Dubrovin, A. T. Fomenko and S. P. Novikov, *Modern Geometry-Methods and Applications*. Springer-Verlag. New York, (1984).
- [20] J. H. Milnor, *Topology from the Differential Viewpoint*. Univ. of Virginia Press, Charlottesville, VA. (1965).
- [21] Y. S. Duan and L. B. Fu, *J. Math. Phys.* **39**, 4343 (1998).
- [22] S. S. Chern, *Ann. Math.* **46**, 674 (1945); **45** 747 (1944).
- [23] Victor Guillemin and Alan Pollack, *Differential Topology*, Prentice-Hall Inc, Englewood Cliffs, New Jersey, 1974.